


Beauseart Soil (loamy sand)



Thorndale Soil (loam)




Fig. 1. Soil pH, # MS germinated out of 50 counted, NH₃ concentration in soil solution, NO₂ and NO₃ content in Beauseart and Thorndale soil amended with various rates of MBM (n=3; ±standard error).




Fig. 2. Number MS germinated out of 50 counted, soil pH, NH₃ concentration in soil solution, NO₃⁻ and NO₂⁻ content of Thorndal soil amended with various rates of MBM (n=3; ±standard error).

004220" 86042960




Fig. 3. Number MS germinated out of 50 counted, NH₃ and HNO₂ concentration in soil solution in Thorndale soil amended with various rates of urea (n=3; \pm standard error).

004220" 86042960




Fig. 4. Number MS germinated out of 50 counted, soil pH, NH₃ and HNO₂ concentration in soil solution in Beauseart soil (sandy loam) amended with various rates of MBM, with and without the nitrification inhibitor, dicyandiamide (DCD) added (n=3; ±standard error).




Fig. 5. Number of MS germinated out of 50 counted and HNO₂ concentration in soil solution of Beauseart soil amended with various fertilizer-N sources to 200 or 400 mg N kg⁻¹ (n=3; ±standard error).

004220"86042960




Fig. 6. Number of MS germinated out of 50 counted, soil pH and HNO₂ concentration in soil solution of Mackenzie soil amended with various amounts of (NH₄)₂SO₄, with and without the nitrification inhibitor, dicyandiamide (DCD) added (n=3; ±standard error).




Fig. 7. Number of MS germinated out of 25 counted exposed for 2 weeks to various concentrations of NH_3 in solid medium ($n=3$; \pm standard error).




Fig. 8. Number of MS germinated out of 50 counted exposed for 8 h, 1 d and 4 d to various concentrations of NH_3 in glycine buffer at pH 8.6 ($n=3$; \pm standard error).

004220-25042360




Fig. 9. Number of MS germinated out of 50 counted exposed for 8 h, 1 d and 4 d to various concentrations of HNO_2 in citric acid buffer at pH 5.0 ($n=3$; \pm standard error).




Fig. 10. Number of MS germinated out of 50 counted after suspension for 4 d to various concentrations of 30 mL HNO_2 citric acid buffer at pH 5.0 in a 250 mL sealer jar ($n=3$; \pm standard error).

004220" 86042960




Fig. 11. Peak concentration of NH_3 in soil solution measured for 12 soils amended with 2% MBM (w w⁻¹) (n=3; \pm standard error).




Fig. 12. Soil pH in response to addition of H_2SO_4 . Soils in Group 1 are in filled symbols and Group 2 are in open symbols.

004220" 86042360




Fig. 13. Number of MS germinated out of 50 counted, soil pH, NO₂⁻ and NO₃⁻ content, and HNO₂ concentration in soil solution of Beauseart and Mackenzie soil amended with (NH₄)₂SO₄ to 200 or 400 mg N kg⁻¹ (n=3).




Fig. 14. Germination of *V. dahliae* MS after submergence in a 0.02 M citric acid buffered solution for a) 1 d exposure to various concentrations of NO_2^- at a solution pH of 4, 5 or 6 and b) 8 h, 1 and 4 d exposure in, or suspended above for 4 d a solution of pH 5 containing various levels of HNO_2 . The concentration of HNO_2 was estimated based on the concentration of NaNO_2 and pH of the solution. Means (± 1 se) of six replicates are shown.

Cont
B8




Fig. 15. Percent colony forming units (cfu) of a) spores of *Streptomyces scabies*, and b) chlamydospores of FOL, and germination of c) sclerotia of *Sclerotinia sclerotiorum*, and seeds of d) *A. retroflexus*, e) *Lutuca sativa* and f) *Raphanus sativus* after submergence for 8 h, 1 and 4 d in 0.02 M citric acid buffered solutions (pH 5.0) containing various levels of HNO_2 . The concentration of HNO_2 was estimated based on the concentration of NaNO_2 and pH of the solution. Means (± 1 se) of six replicates are shown.